Author Index

Aanesrud, A., see Iversen, B. 217
Arimoto, H., see Kawano, H. 71
Aronson, A.L., see Tyczkowska, K. 337
Aucoin, D.P., see Tyczkowska, K. 337
Bagchi, S.P., see Jindal, S.P. 392
Bartlett, W.A.
Effects of mobile phase composition on the chromatographic and electrochemical behaviour of catecholamines and selected metabolites. Reversed-phase ion-paired high-performance liquid chromatography using multiple-electrode detection 1
Benet, L.Z., see Teng, R.-L. 421
Blair, I.A., see Prakash, C. 325
Bochner, F., see Chen, Z.R. 313
Boldizsár, M., see Kremmer, T. 45
Breckenridge, A.M., see Idowu, O.R. 125
Brendel, E., see Meineke, I. 287
Brusilow, S.W.
— and Hauser, E.
Simple method of measurement of orotic acid and orotidine in urine 388
Bruyere, Jr., H.J., see Kapil, R.P. 182
Carlier, J.-P.
— and Sellier, N.
Gas chromatographic–mass spectral studies after methylation of metabolites produced by some anaerobic bacteria in spent media 257
Chen, Z.R.
—, Reynolds, G., Bochner, F. and Somogyi, A.
Direct determination of codeine-6-glucuronide in plasma and urine using solid-phase extraction and high-performance liquid chromatography with fluorescence detection 313
Cheymol, G., see Poirier, J.-M. 409
Clark, C.R., see Lukey, B.J. 117
Costabel, U., see Maier, K. 380
Dehelean, L., see Selinger, K. 230
De Mey, C., see Meineke, I. 287
De Vries, J.X.
— and Völker, U.
Separation of the enantiomers of phenprocoumon and warfarin by high-performance liquid chromatography using a chiral stationary phase. Determination of the enantiomeric ratio of phenprocoumon in human plasma and urine 149
Edwards, G., see Idowu, O.R. 125
Eisenberg, E.J.
—, Patterson, W.R. and Kahn, G.C.
High-performance liquid chromatographic method for the simultaneous determination of the enantiomers of carvedilol and its O-desmetyl metabolite in human plasma after chiral derivatization 105
Fay, L., see Michel, G. 188
Febles, M., see Tápenes, R.D. 202
Filser, J.G.
—, Koch, S., Fisher, M. and Müller, W.E.
Determination of urinary 3-methoxy-4-hydroxyphenylethylene glycol and its conjugates by high-performance liquid chromatography with electrochemical and ultraviolet absorbance detection 275
Fisher, M., see Filser, J.G. 275
Fujii, T., see Kawano, H. 71
Fukasawa, K., see Harada, M. 176
Fukasawa, K.M., see Harada, M. 176
Garner, D.M., see Pieniaszek, Jr., H.J. 79
Goto, J.
—, Miura, H. and Nambara, T.
Studies on steroids. CCXXXXV. Determination of 5β-cholestanoic acids in human urine by gas chromatography–mass spectrometry with negative ion chemical ionization detection 245
Guerrieri, A., see Palmisano, F. 35
Harada, M.
—, Mogi, M., Fukasawa, K. and Fukasawa, K.M.
High-performance liquid chromatographic determination of aminopeptidase P activity in Fischer F344 rat serum and kidney 176
Hauser, E., see Brusilow, S.W. 388
Hedeen, K.M., see Tyczkowska, K. 337
Henne, E.M., see Meineke, I. 287
Herrmann, T., see Steinhilber, D. 361
Hill, H.M., see Selinger, K. 230
Ho, Y., see Hsu, K.-Y. 305
Holényi, I., see Kremmer, T. 45
Hsu, K.-Y.
— and Ho, Y.
Determination of isoniazid methanesulphonate and its metabolites in rabbit blood

by high-performance liquid chromatography 305
Idowu, O.R.
—, Edwards, G., Ward, S.A., Orme, M.L'E. and Breckenridge, A.M.
Determination of arteether in blood plasma by high-performance liquid chromatography with ultraviolet detection after hydrolysis with acid 125
Igaki, A.
—, Kobayashi, K., Kimura, M., Sakoguchi, T. and Matsuoka, A.
Determination of serum sulphonylureas by high-performance liquid chromatography with fluorimetric detection 222
Inotsume, N., see Kawano, H. 71
Isensee, R.K., see Pieniaszek, Jr., H.J. 79
Ishizaki, T., see Kubota, K. 373
Ito, K., see Sato, J. 239
Iversen, B.
—, Aanesrud, A., Kolstad, A.K. and Rasmussen, K.E.
Determination of oxytetracycline in plasma from rainbow trout using high-performance liquid chromatography with ultraviolet detection 217
Izawa, K., see Yagi, S. 27
Jajoo, H.K., see Prakash, C. 325
Jindal, S.P.
—, Lutz, T. and Bagchi, S.P.
Mass spectrometric determination of tetrabenazine using a stable isotope-labeled analogue as an internal standard 392
Kahn, G.C., see Eisenberg, E.J. 105
Kapil, R.P.
— and Bruyere, Jr., H.J.
Applicability of capillary gas chromatography to study caffeine distribution in the developing chicken egg 182
Karnes, H.T., see Mallikaarjun, S. 93
Kato, H., see Moriyama, T. 164
Kawano, H.
—, Inotsume, N., Arimoto, H., Fujii, T. and Nakano, M.
Rapid and sensitive determination of aprindine in serum by gas chromatography using a surface ionization detector 71
Kennedy, S.W.
— and Maslen, A.L.
Separation of porphyrin isomers by high-performance liquid chromatography 53
Kim, O., see Shin, H. 398
Kimura, M., see Igaki, A. 222
Kobayashi, K., see Igaki, A. 222
Kobayashi, T., see Okano, T. 63
Koch, S., see Filser, J.G. 275
Kolstad, A.K., see Iversen, B. 217
Krämer, B.K.
—, Ress, K.M., Mayer, F., Kühlkamp, V., Liebich, H.M., Risler, T. and Seipel, L.
Rapid high-performance liquid chromatographic method for the quantification of mexiletine and its metabolites in serum 414
Kremmer, T.
—, Paulik, E., Boldizsár, M. and Holényi, I.
Application of the fast protein liquid chromatographic system and MonoQ HR 5/5 anion exchanger to the separation of nucleotides 45
Kubota, K.
—, Ishizaki, T. and Oka, K.
Chromatographic determination of percutaneous absorption of topical non-radiolabelled prednisolone in vivo, and preliminary application to transdermal pharmacokinetics 373
Kühlkamp, V., see Krämer, B.K. 414
Kunze, D., see Rabe, H. 353
Lebot, M., see Poirier, J.-M. 409
Leff, A.R., see Munoz, N.M. 157
Lemonnier, A., see Myara, I. 170
Lenz, A.-G., see Maier, K. 380
Leroy, P., see Liu, H.-F. 137
Leuschel, L., see Maier, K. 380
Lho, D., see Shin, H. 398
Liebich, H.M., see Krämer, B.K. 414
Lieske, C.N., see Lukey, B.J. 117
Liu, H.-F.
—, Leroy, P., Nicolas, A., Magdalou, J. and Siest, G.
Evaluation of a versatile reversed-phase high-performance liquid chromatographic system using cethexonium bromide as ion-pairing reagent for the analysis of glucuronic acid conjugates 137
Lukey, B.J.
—, Marlow, D.D., Clark, C.R., McCluskey, M.P. and Lieske, C.N.
Application of a new radiometric high-performance liquid chromatographic as-

say to define physostigmine pharmacokinetics in guinea pigs 117
Lutz, T., see Jindal, S.P. 392
McCluskey, M.P., see Lukey, B.J. 117
Magdalou, J., see Liu, H.-F. 137
Maier, K.
—, Costabel, U., Lenz, A.-G. and Leuschel, L.
Simultaneous determination of L-homoserine and L-homoserine lactone by reversed-phase liquid chromatography in acid hydrolysates of proteins after cyanogen bromide treatment 380
Makino, I., see Moriyama, T. 164
Makino, M., see Moriyama, T. 164
Mallikaarjun, S.
—, Wood, J.H. and Karnes, H.T.
High-performance liquid chromatographic method for the determination of salicylic acid and its metabolites in urine by direct injection 93
Marlow, D.D., see Lukey, B.J. 117
Maslen, A.L., see Kennedy, S.W. 53
Matheou, D., see Selinger, K. 230
Matsuoka, A., see Igaki, A. 222
Matsuura, S., see Niwa, T. 347
Mayer, F., see Krämer, B.K. 414
Mayol, R.F., see Prakash, C. 325
Mehvar, R.
Stereospecific liquid chromatographic analysis of racemic adrenergic drugs utilizing precolumn derivatization with (−)-menthyl chloroformate 402
Meineke, I.
—, Stüwe, E., Henne, E.M., Rusteberg, G., Brendel, E. and De Mey, C.
Routine measurement of plasma catecholamines in clinical pharmacology by high-performance liquid chromatography with electrochemical detection 287
Mey, C. De, see Meineke, I. 287
Michel, G.
—, Fay, L. and Prost, M.
Determination of isosorbide dinitrate and its metabolites in plasma by gas chromatography on a capillary column with electron-capture detection 188
Miura, H., see Goto, J. 245
Moatti, N., see Myara, I. 170
Mogi, M., see Harada, M. 176
Mohri, Z.-I., see Yagi, S. 27
Moriyama, T.
—, Yamadera, K., Takebe, T., Makino, I., Kato, H., Nobuoka, M. and Makino, M.
Purification of the pancreatic stone protein by high-performance liquid chromatography 164
Müller, W.E., see Filser, J.G. 275
Munoz, N.M.
—, Tutins, C. and Leff, A.R.
Highly sensitive determination of catecholamine and serotonin concentrations in plasma by liquid chromatography–electrochemistry 157
Myara, I.
—, Moatti, N. and Lemonnier, A.
Separation of two erythrocyte prolidase isoforms by fast protein liquid chromatography: application to prolidase deficiency 170
Nagatsu, T., see Niwa, T. 347
Nakagawa, T., see Yagi, S. 27
Nakagawa, Y., see Rabe, H. 353
Nakano, M., see Kawano, H. 71
Nambara, T., see Goto, J. 245
Nicolas, A., see Liu, H.-F. 137
Niida, Y., see Sato, J. 239
Niwa, T.
—, Yoshizumi, H., Tatematsu, A., Matsuura, S. and Nagatsu, T.
Presence of tetrahydroisoquinoline, a parkinsonism-related compound, in foods 347
Nobuoka, M., see Moriyama, T. 164
Nohta, H.
—, Yamaguchi, E., Ohkura, Y. and Watanabe, H.
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by high-performance liquid chromatography with fluorescence derivatization 15
Ohkura, Y., see Nohta, H. 15
Oka, K., see Kubota, K. 373
Okano, T.
—, Tsuwaga, N. and Kobayashi, T.
Identification and determination of 1α,25-dihydroxyvitamin D_3 in rat skin by high-performance liquid chromatography and radioreceptor assay 63

Orme, M.L'E., see Idowu, O.R. 125
Owada, E., see Sato, J. 239
Palmisano, F.
—, Rotunno, R., Guerrieri, A. and Zambonin, P.G.
Simultaneous determination of pseudouridine and creatinine in untreated urine by ion-pair liquid chromatography with diode-array ultraviolet detection 35
Paris, A.
—, Sutra, J.F. and Rao, D.
Separation of C-17 fatty acid of 17β-estradiol by reversed-phase high-performance liquid chromatography 367
Park, J., see Shin, H. 398
Patterson, W.R., see Eisenberg, E.J. 105
Paulik, E., see Kremmer, T. 45
Pérez Avila, J., see Tápanes, R.D. 196, 202
Pieniaszek, Jr., H.J.
—, Shen, H.-S.L., Garner, D.M., Page, G.O., Shalaby, L.M., Isensee, R.K. and Whitney, Jr., C.C.
Determination of unlabeled and $^{13}C_6$-labeled moricizine in human plasma using thermospray liquid chromatography–mass spectrometry 79
Poirier, J.-M.
—, Lebot, M. and Cheymol, G.
Rapid and sensitive column liquid chromatographic determination of sotalol in plasma 409
Prakash, C.
—, Jajoo, H.K., Blair, I.A. and Mayol, R.F.
Resolution of enantiomers of the antiarrhythmic drug encainide and its major metabolites by chiral derivatization and high-performance liquid chromatography 325
Prost, M., see Michel, G. 188
Rabe, H.
—, Reichmann, G., Nakagawa, Y., Rüstow, B. and Kunze, D.
Separation of alkylacyl and diacyl glycerophospholipids and their molecular species as naphthylurethanes by high-performance liquid chromatography 353
Ramachandran, P.K., see Rao, S.S. 210
Ramos, F., see Tápanes, R.D. 196, 202
Rao, D., see Paris, A. 367
Rao, S.S.
—, Vijayaraghavan, R., Suryanarayana, M.V.S. and Ramachandran, P.K.
Gas chromatographic identification of urinary metabolites of insect repellent N,N-diethylphenylacetamide on inhalation exposure in rats 210
Rasmussen, K.E., see Iversen, B. 217
Reichmann, G., see Rabe, H. 353
Ress, K.M., see Krämer, B.K. 414
Reynolds, G., see Chen, Z.R. 313
Risler, T., see Krämer, B.K. 414
Roth, H.J., see Steinhilber, D. 361
Rotunno, R., see Palmisano, F. 35
Rüstow, B., see Rabe, H. 353
Rusteberg, G., see Meineke, I. 287
Sakoguchi, T., see Igaki, A. 222
Sato, J.
—, Owado, E., Ito, K., Niida, Y., Wakamatsu, A. and Umetsu, M.
Simple, rapid and sensitive reversed-phase high-performance liquid chromatographic method for the determination of mefenamic acid in plasma 239
Seipel, L., see Krämer, B.K. 414
Selinger, K.
—, Hill, H.M., Matheou, D. and Dehelean, L.
Determination of free and total metaproterenol in human plasma by high-performance liquid chromatography with fluorimetric detection 230
Sellier, N., see Carlier, J.-P. 257
Shalaby, L.M., see Pieniaszek, Jr., H.J. 79
Shen, H.-S.L., see Pieniaszek, Jr., H.J. 79
Shin, H.
—, Park, J., Lho, D. and Kim, O.
Measurement of zipeprol in rat plasma by gas chromatography with nitrogen–phosphorus detection 398
Siest, G., see Liu, H.-F. 137
Somogyi, A., see Chen, Z.R. 313
Steinhilber, D.
—, Herrmann, T. and Roth, H.J.
Separation of lipoxins and leukotrienes from human granulocytes by high-performance liquid chromatography with a Radial-Pak cartridge after extraction with an octadecyl reversed-phase column 361
Stüwe, E., see Meineke, I. 287
Suryanarayana, M.V.S., see Rao, S.S. 210
Sutra, J.F., see Paris, A. 367
Takebe, T., see Moriyama, T. 164

Tanaka, H., see Yagi, S. 27
Tápanes, R.D.
—, Ramos, F., Febles, M. and Pérez Avila, J.
Determination of dabequin in biological fluids by gas chromatography with a nitrogen-selective detector 202
—, Ramos, F. and Pérez Avila, J.
Contribution to the determination of chloroquine in biological fluids 196
Tatematsu, A., see Niwa, T. 347
Teng, R.-L.
— and Benet, L.Z.
Simultaneous measurement of prednisone and 6β-hydroxyprednisolone in urine by high-performance liquid chromatography using dexamethasone as the internal standard 421
Tsugawa, N., see Okano, T. 63
Tutins, C., see Munoz, N.M. 157
Tyczkowska, K.
—, Hedeen, K.M., Aucoin, D.P. and Aronson, A.L.
High-performance liquid chromatographic method for the simultaneous determination of enrofloxacin and its primary metabolite ciprofloxacin in canine serum and prostatic tissue 337
Umetsu, M., see Sato, J. 239
Vijayaraghavan, R., see Rao, S.S. 210
Völker, U., see De Vries, J.X. 149
Vries, J.X. De, see De Vries, J.X. 149
Wakamatsu, A., see Sato, J. 239
Ward, S.A., see Idowu, O.R. 125
Watanabe, H., see Nohta, H. 15
Whitney, Jr., C.C., see Pieniaszek, Jr., H.J. 79
Wood, J.H., see Mallikaarjun, S. 93
Yagi, S.
—, Izawa, K., Nakagawa, T., Tanaka, H., Yoshitake, A. and Mohri, Z.-I.
Efficient high-performance liquid chromatographic system for protein purification 27
Yamadera, K., see Moriyama, T. 164
Yamaguchi, E., see Nohta, H. 15
Yoshitake, A., see Yagi, S. 27
Yoshizumi, H., see Niwa, T. 347
Zambonin, P.G., see Palmisano, F. 35

Subject Index

Acetylisoniazid
Determination of isoniazid methanesulphonate and its metabolites in rabbit blood by HPLC 305

Adrenergic drugs, racemic
Stereospecific LC analysis of racemic adrenergic drugs utilizing precolumn derivatization with (−)-menthyl chloroformate 402

Alkylacylglycerol
Separation of alkylacyl and diacyl glycerophospholipids and their molecular species as naphthylurethanes by HPLC 353

L-Amino acids
Simultaneous determination of L-homoserine and L-homoserine lactone by reversed-phase LC in acid hydrolysates of proteins after cyanogen bromide treatment 380

Aminopeptidase P
HPLC determination of aminopeptidase P activity in Fischer F344 rat serum and kidney 176

Aprindine
Rapid and sensitive determination of aprindine in serum by GC using a surface ionization detector 71

Arteether
Determination of arteether in blood plasma by HPLC with UV detection after hydrolysis with acid 125

Atenolol
Stereospecific LC analysis of racemic adrenergic drugs utilizing precolumn derivatization with (−)-menthyl chloroformate 402

Caffeine
Applicability of capillary GC to study caffeine distribution in the developing chicken egg 182

Carvedilol
HPLC method for the simultaneous determination of carvedilol and its O-desmethyl metabolite in human plasma after chiral derivatization 105

Catecholamines
Effects of mobile phase composition on the chromatographic and electrochemical behaviour of catecholamines and selected metabolites. Reversed-phase ion-paired HPLC using multiple-electrode detection 1

Catecholamines
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15

Catecholamines
Highly sensitive determination of catecholamine and serotonin concentrations in plasma by LC–electrochemistry 157

Catecholamines
Routine measurement of plasma catecholamines in clinical pharmacology by HPLC with electrochemical detection 287

Cethexonium bromide
Evaluation of a versatile reversed-phase HPLC system using cethexonium bromide as ion-pairing reagent for the analysis of glucuronic acid conjugates 137

Chloroquine
Contribution to the determination of chloroquine in biological fluids 196

5β-Cholestanoic acids
Studies on steroids. CCXXXXV. Determination of 5β-cholestanoic acids in human urine by GC–MS with negative ion chemical ionization detection 245

Ciprofloxacin
HPLC method for the simultaneous determination of enrofloxacin and its primary metabolite ciprofloxacin in canine serum and prostatic tissue 337

Clofibric acid
Evaluation of a versatile reversed-phase HPLC system using cethexonium bromide as ion-pairing reagent for the analysis of glucuronic acid conjugates 137

Codeine-6-glucuronide
Direct determination of codeine-6-glucuronide in plasma and urine using solid-phase extraction and HPLC with fluorescence detection 313

Coproporphyrins
Separation of porphyrin isomers by HPLC 53
Creatinine
Simultaneous determination of pseudouridine and creatinine in untreated urine by ion-pair LC with diode-array UV detection 35
Dabequin
Determination of dabequin in biological fluids by GC with a nitrogen-selective detector 202
N-Demethylencainide
Resolutions of enantiomers of the antiarrhythmic drug encainide and its major metabolites by chiral derivatization and HPLC 325
O-Demethylencainide
Resolutions of enantiomers of the antiarrhythmic drug encainide and its major metabolites by chiral derivatization and HPLC 325
O-Desmethylcarvedilol
HPLC method for the simultaneous determination of carvedilol and its O-desmethyl metabolite in human plasma after chiral derivatization 105
Diacylglycerol
Separation of alkylacyl and diacyl glycerophospholipids and their molecular species as naphthylurethanes by HPLC 353
N,N-Diethylphenylacetamide
GC identification of urinary metabolites of insect repellent N,N-diethylphenylacetamide on inhalation exposure in rats 210
3,4-Dihydroxybenzylamine
Effects of mobile phase composition on the chromatographic and electrochemical behaviour of catecholamines and selected metabolites. Reversed-phase ion-paired HPLC using multiple-electrode detection 1
3α,7α-Dihydroxy-5β-cholestanoic acid
Studies on steroids. CCXXXXV. Determination of 5β-cholestanoic acids in human urine by GC–MS with negative ion chemical ionization detection 245
3,4-Dihydroxymandelic acid
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15
3,4-Dihydroxyphenylacetic acid
Effects of mobile phase composition on the chromatographic and electrochemical behaviour of catecholamines and selected metabolites. Reversed-phase ion-paired HPLC using multiple-electrode detection 1
3,4-Dihydroxyphenylacetic acid
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15
3,4-Dihydroxyphenylalanine
Effects of mobile phase composition on the chromatographic and electrochemical behaviour of catecholamines and selected metabolites. Reversed-phase ion-paired HPLC using multiple-electrode detection 1
3,4-Dihydroxyphenylethylene glycol
Effects of mobile phase composition on the chromatographic and electrochemical behaviour of catecholamines and selected metabolites. Reversed-phase ion-paired HPLC using multiple-electrode detection 1
3,4-Dihydroxyphenylethylene glycol
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15
1α,25-Dihydroxyvitamin D_3
Identification and determination of 1α,25-dihydroxyvitamin D_3 in rat skin by HPLC and radioreceptor assay 63
L-DOPA
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15
Dopamine
Effects of mobile phase composition on the chromatographic and electrochemical behaviour of catecholamines and selected metabolites. Reversed-phase ion-paired

HPLC using multiple-electrode detection 1

Dopamine
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15

Dopamine
Routine measurement of plasma catecholamines in clinical pharmacology by HPLC with electrochemical detection 287

Encainide
Resolution of enantiomers of the antiarrhythmic drug encainide and its major metabolites by chiral derivatization and HPLC 325

Enrofloxacin
HPLC method for the simultaneous determination of enrofloxacin and its primary metabolite ciprofloxacin in canine serum and prostatic tissue 337

Enzymes
Separation of two erythrocyte prolidase isoforms by FPLC; application to prolidase deficiency 170

Enzymes
HPLC determination of aminopeptidase P activity in Fischer F344 rat serum and kidney 176

Epinephrine
Effects of mobile phase composition on the chromatographic and electrochemical behaviour of catecholamines and selected metabolites. Reversed-phase ion-paired HPLC using multiple-electrode detection 1

Epinephrine
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15

Epinephrine
Highly sensitive determination of catecholamine and serotonin concentrations in plasma by LC–electrochemistry 157

Epinephrine
Routine measurement of plasma catecholamines in clinical pharmacology by HPLC with electrochemical detection 287

17β-Estradiol, C-17 fatty acid esters
Separation of C-17 fatty acid esters of 17β-estradiol by reversed-phase HPLC 367

Fatty acids
GC–MS studies after methylation of metabolites produced by some anaerobic bacteria in spent media 257

Gentisic acid
HPLC method for the determination of salicylic acid and its metabolites in urine by direct injection 93

Glucuronic acid conjugates
Evaluation of a versatile reversed-phase HPLC system using cethexonium bromide as ion-pairing reagent for the analysis of glucuronic acid conjugates 137

Glycerophospholipids
Separation of alkylacyl and diacyl glycerophospholipids and their molecular species as naphthylurethanes by HPLC 353

Growth hormone receptors
Efficient HPLC system for protein purification 27

L-Homoserine
Simultaneous determination of L-homoserine and L-homoserine lactone by reversed-phase LC in acid hydrolysates of proteins after cyanogen bromide treatment 380

L-Homoserine lactone
Simultaneous determination of L-homoserine and L-homoserine lactone by reversed-phase LC in acid hydrolysates of proteins after cyanogen bromide treatment 380

Homovanillic acid
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15

4-Hydroxy-3-methoxyphenylethylene glycol
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15

6β-Hydroxyprednisolone
Simultaneous measurement of prednisone, prednisolone and 6β-hydroxypred-

nisolone in urine by HPLC using dexamethasone as the internal standard 421
Isoniazid
Determination of isoniazid methanesulphonate and its metabolites in rabbit blood by HPLC 305
Isoniazid methanesulphonate
Determination of isoniazid methanesulphonate and its metabolites in rabbit blood by HPLC 305
1-Isopropylamino-3-phenoxypropan-2-ol
Stereospecific LC analysis of racemic adrenergic drugs utilizing precolumn derivatization with (−)-menthyl chloroformate 402
Isosorbide dinitrate
Determination of isosorbide dinitrate and its metabolites in plasma by GC on a capillary column with electron-capture detection 188
Isosorbide mononitrates
Determination of isosorbide dinitrate and its metabolites in plasma by GC on a capillary column with electron-capture detection 188
Leukotrienes
Separation of lipoxins and leukotrienes from human granulocytes by HPLC with a Radial-Pak cartridge after extraction with an octadecyl reversed-phase column 361
Lipoxins
Separation of lipoxins and leukotrienes from human granulocytes by HPLC with a Radial-pak cartridge after extraction with an octadecyl reversed-phase column 361
Mefenamic acid
Simple, rapid and sensitive reversed-phase HPLC method for the determination of mefenamic acid in plasma 239
Mesoporphyrin III
Separation of porphyrin isomers by HPLC 53
Metanephrine
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15
Metaproterenol
Determination of free and total metaproterenol in human plasma by HPLC with fluorimetric detection 230
Metaproterenol
Stereospecific LC analysis of racemic adrenergic drugs utilizing precolumn derivatization with (−)-menthyl chloroformate 402
3-Methoxy-O-demethylencainide
Resolution of enantiomers of the antiarrhythmic drug encainide and its major metabolites by chiral derivatization and HPLC 325
3-Methoxy-4-hydroxyphenylethylene glycol
Determination of urinary 3-methoxy-4-hydroxyphenylethylene glycol and its conjugates by HPLC with electrochemical and UV absorbance detection 275
3-Methoxytyramine
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15
Metoprolol
Stereospecific LC analysis of racemic adrenergic drugs utilizing precolumn derivatization with (−)-menthyl chloroformate 402
Mexiletine
Rapid HPLC method for the quantification of mexiletine and its metabolites in serum 414
Mobile phase conditions
Effects of mobile phase composition on the chromatographic and electrochemical behaviour of catecholamines and selected metabolites. Reversed-phase ion-paired HPLC using multiple-electrode detection 1
Moricizine
Determination of unlabeled and $^{13}C_6$-labeled moricizine in human plasma using thermospray LC–MS 79
Non-volatile fatty acids
GC–MS studies after methylation of metabolites produced by some anaerobic bacteria in spent media 257
Norepinephrine
Effects of mobile phase composition on the chromatographic and electrochemical be-

haviour of catecholamines and selected metabolites. Reversed-phase ion-paired HPLC using multiple-electrode detection 1

Norepinephrine
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15

Norepinephrine
Highly sensitive determination of catecholamine and serotonin concentrations in plasma by LC–electrochemistry 157

Norepinephrine
Routine measurement of plasma catecholamines in clinical pharmacology by HPLC with electrochemical detection 287

Normetanephrine
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15

Nucleotides
Application of the FPLC system and MonoQ HR 5/5 anion exchanger to the separation of nucleotides 45

Orotic acid
Simple method of measurement of orotic acid and orotidine in urine 388

Orotidine
Simple method of measurement of orotic acid and orotidine in urine 388

Oxytetracycline
Determination of oxytetracycline in plasma from rainbow trout using HPLC with UV detection 217

Pancreatic stone protein
Purification of the pancreatic stone protein by HPLC 164

Phenprocoumon
Separation of the enantiomers of phenprocoumon and warfarin by HPLC using a chiral stationary phase. Determination of the enantiomeric ratio of phenprocoumon in human plasma and urine 149

Physostigmine
Application of a new radiometric HPLC assay to define physostigmine pharmacokinetics in guinea pigs 117

Porphyrins
Separation of porphyrin isomers by HPLC 53

Prednisolone
Chromatographic determination of percutaneous absorption of topical non-radiolabelled prednisolone in vivo, and preliminary application to transdermal pharmacokinetics 373

Prednisolone
Simultaneous measurement of prednisone, prednisolone and 6β-hydroxyprednisolone in urine by HPLC using dexamethasone as the internal standard 421

Prednisone
Simultaneous measurement of prednisone, prednisolone and 6β-hydroxyprednisolone by HPLC using dexamethasone as the internal standard 421

Prolidases
Separation of two erythrocyte prolidase isoforms by FPLC; application to prolidase deficiency 170

Propranolol
Stereospecific LC analysis of racemic adrenergic drugs utilizing precolumn derivatization with (−)-menthyl chloroformate 402

Proteins
Efficient HPLC system for protein purification 27

Proteins
Purification of the pancreatic stone protein by HPLC 164

Pseudouridine
Simultaneous determination of pseudouridine and creatinine in untreated urine by ion-pair LC with diode-array UV detection 35

Salicyl acyl glucuronide
HPLC method for the determination of salicylic acid and its metabolites in urine by direct injection 93

Salicylhydroxamic acid
HPLC method for the determination of salicylic acid and its metabolites in urine by direct injection 93

Salicylic acid
HPLC method for the determination of salicylic acid and its metabolites in urine by direct injection 93

Salicyl phenolic glucuronide
HPLC method for the determination of salicylic acid and its metabolites in urine by direct injection 93
Salicyluric acid
HPLC method for the determination of salicylic acid and its metabolites in urine by direct injection 93
Serotonin
High sensitive determination of catecholamine and serotonin concentrations in plasma by LC–electrochemistry 157
Sotalol
Stereospecific LC analysis of racemic adrenergic drugs utilizing precolumn derivatization with (−)-menthyl chloroformate 402
Sotalol
Rapid and sensitive column LC determination of sotalol in plasma 409
Steroids
Studies on steroids. CCXXXXV. Determination of 5β-cholestanoic acids in human urine by GC–MS with negative ion chemical ionization detection 245
Steroids
Separation of C-17 fatty acid esters of 17β-estradiol by reversed-phase HPLC 367
Steroids
Chromatographic determination of percutaneous absorption of topical non-radiolabelled prednisolone in vivo, and preliminary application to transdermal pharmacokinetics 373
Steroids
Simultaneous measurement of prednisone, prednisolone and 6β-hydroxyprednisolone in urine by HPLC using dexamethasone as the internal standard 421
Sulphonylureas
Determination of serum sulphonylureas by HPLC with fluorimetric detection 222
Tetrabenazine
MS determination of tetrabenazine using a stable isotope-labeled analogue as an internal standard 392
Tetrahydroisoquinoline
Presence of tetrahydroisoquinoline, a parkinsonism-related compound, in foods 347
Toliprolol
Stereospecific LC analysis of racemic adrenergic drugs utilizing precolumn derivatization with (−)-menthyl chloroformate 402
3α,7α,12α-Trihydroxy-5β-cholestanoic acid
Studies on steroids. CCXXXXV. Determination of 5β-cholestanoic acids in human urine by GC–MS with negative ion chemical ionization detection 245
Uroporphyrins
Separation of porphyrin isomers by HPLC 53
Vanillylmandelic acid
Measurement of catecholamines, their precursor and metabolites in human urine and plasma by solid-phase extraction followed by HPLC with fluorescence derivatization 15
Volatile fatty acids
GC–MS studies after methylation of metabolites produced by some anaerobic bacteria in spent media 257
Warfarin
Separation of the enantiomers of phenprocoumon and warfarin by HPLC using a chiral stationary phase. Determination of the enantiomeric ratio of phenprocoumon in human plasma and urine 149
Zipeprol
Measurement of zipeprol in rat plasma by GC with nitrogen–phosphorus detection 398